arXiv:1611.06570v2 [math.CO] 17 Dec 2016

Square of a Hamilton cycle in a random graph

Patrick Bennett* Andrzej Dudek† Alan Frieze‡

December 20, 2016

Abstract

We show that the threshold for the random graph $G_{n,p}$ to contain the square of a Hamilton cycle is $p = \frac{1}{\sqrt{n}}$. This improves the previous results of Kühn and Osthus and also Nenadov and Škorić. In addition we consider how many random edges need to be added to a graph of order n with minimum degree αn in order that it contains the square of a Hamilton cycle w.h.p.

1 Introduction

Let us recall a few classic models of random graphs. The *random graph* $G_{n,m}$ is the random graph G with vertex set $[n]$ and m random edges. The *random graph* $G_{n,p}$ is the random graph G with vertex set $[n]$ in which each of the $\binom{n}{2}$ edges of the complete graph K_n is included independently with probability p. Similarly, the *random digraph* $D_{n,m}$ is the random digraph D with vertex set $[n]$ and m random edges. The *random digraph* $D_{n,p}$ is the random digraph D with vertex set $[n]$ in which each of the $n(n-1)$ edges of the complete digraph $\vec{K}_n$ is included independently with probability p. We say that a sequence of events $\mathcal{E}_n$ in a probability space holds *with high probability* (or *w.h.p.*) if the probability that $\mathcal{E}_n$ holds tends to 1 as $n \to \infty$. Throughout this note all logarithms are natural (base e) and all asymptotics are taken in n.

By the *kth power of a Hamilton cycle*, we mean a permutation (bijection) $\pi : [n] \to [n]$ such that $\{\pi(i), \pi(j)\} \in E(G)$ whenever $i < j \leq i + k$. (Here $i + k$ is to be taken as $i + k - n$ if $i + k \geq n + 1$.) Hamilton cycles have long been studied in the context of random graphs (see, e.g., [1, 3, 7, 13]). Powers of Hamilton cycles are less well-studied and much less is known about

*Department of Mathematics, Department of Mathematics, Western Michigan University, Kalamazoo, MI. Supported in part by Simons Foundation Grant #426894.

†Department of Mathematics, Department of Mathematics, Western Michigan University, Kalamazoo, MI. Supported in part by Simons Foundation Grant #244712 and by the National Security Agency under Grant Number H98230-15-1-0172. The United States Government is authorized to reproduce and distribute reprints notwithstanding any copyright notation hereon.

‡Department of Mathematical Sciences, Carnegie Mellon University, Pittsburgh PA. Research supported in part by NSF grant CCF1013110 and Simons Foundation Grant #1030891.

them. Since the kth power of a Hamilton cycle contains kn edges, we can see that if X_k denotes the number of copies of such in $G_{n,p}$, then by using Stirling's formula,

$$\mathbf{E}(X_k) = n!p^{kn} \sim \sqrt{2\pi n}\left(\frac{n}{e}\right)^n p^{kn} = \sqrt{2\pi n}\left(\frac{np^k}{e}\right)^n$$

and so $\mathbf{E}(X_k) \to 0$ if $p \le \left(\frac{(1-\varepsilon)e}{n}\right)^{1/k}$ for any constant $\varepsilon > 0$. Thus, if $p \le \left(\frac{(1-\varepsilon)e}{n}\right)^{1/k}$, then w.h.p. $G_{n,p}$ contains no kth power of a Hamilton cycle.

Kühn and Osthus [9] observed that for $k \ge 3$, $p = \frac{1}{n^{1/k}}$ is the coarse threshold for the existence of the kth power of a Hamilton cycle in $G_{n,p}$. This comes directly from a result of Riordan [14]. For $k = 2$ they gave a bound of $p \ge n^{-1/2+\varepsilon}$ (for any $\varepsilon > 0$) being sufficient for the existence of the square of a Hamilton cycle w.h.p.. This result was improved by Nenadov and Škorić [11] to $p \ge \frac{C\log^4 n}{\sqrt{n}}$ (C is a positive constant) being sufficient for the existence of the square of a Hamilton cycle. Here we improve this result by decreasing the lower bound to $\frac{\omega}{\sqrt{n}}$, where $\omega \to \infty$. As a matter of fact we prove a stronger result in the $D_{n,m}$ model.

Theorem 1 *If $\omega \to \infty$ and $m = \omega n^{3/2}$, then w.h.p. $D_{n,m}$ contains the square of a Hamilton cycle.*

This theorem also implies that the same holds in $D_{n,p}$, $G_{n,m}$ and $G_{n,p}$ with $p = \frac{\omega}{\sqrt{n}}$. We will give details of this implication at the end of Section 2. In the argument, we replace ω by monotone functions of ω, which is clearly acceptable.

The proof is based on the second moment method. Let us emphasize that applying the second moment method to $G_{n,p}$ or $D_{n,p}$ fails.

In this paper we also consider a problem related to the Posá-Seymour conjecture, which states that every graph G on n vertices with minimum degree at least $kn/(k+1)$ contains the kth power of a Hamilton cycle. This conjecture was proved for large enough n by Komlós, Sarközy and Szemerédi [8]. Bohman, Frieze and Martin [2] considered the question of how many random edges need to be added to a dense graph in order that it is Hamiltonian w.h.p. The following theorem extends their result to the square of a Hamilton cycle. For graphs $G = (V, E)$ and $X = (V, F)$ we define a graph $G + X$ on vertex set V with edge set $E \cup F$.

Theorem 2 *Let G be a graph of order n which has minimum degree at least αn for some absolute constant $\alpha > 1/2$. Let X denote a set of randomly chosen edges. Then w.h.p. $G + X$ contains the square of a Hamilton cycle, provided that*

$$|X| \ge Kn^{4/3}\log^{1/3} n.$$

Here we need $K = K(\alpha)$ to be sufficiently large.

Clearly $n^{4/3}$ is less than the $n^{3/2}$ needed if all edges are random.

2 Outline proof of Theorem 1

We will consider a slightly different model $D^*_{n,2m}$. We generate $D^*_{n,2m}$, where $m = \omega n^{3/2}$, by first choosing m random directed red edges and then choosing m random directed blue edges that are independent of the red edges (so an edge may be both red and blue). This is not a problem for the analysis and no edge will appear twice in the square of a Hamilton cycle. Also, after removing the colors, we see that we have a multigraph. A simple calculation shows that w.h.p. we have $O(\omega^2 n)$ repeated edges and we will assume that ω grows slowly. Thus we will have proved a property of $D_{n,2m-O(\omega^2 n)}$.

We say that a permutation π of $[n]$ in $D^*_{n,2m}$ is *square inducing* if $D^*_{n,2m}$ contains red edges $(\pi(i), \pi(i+1))$ and blue edges $(\pi(i), \pi(i+2))$ for each $i \in [n]$.

We let $\mathcal{H}_\pi$ be the event that permutation π is square inducing in $D^*_{n,2m}$. Let X be the random variable that counts the number of square inducing permutations.

$$\mathbf{E}(X) = \sum_\pi \mathbf{Pr}(\mathcal{H}_\pi) = n! \left(\frac{\binom{N-n}{m-n}}{\binom{N}{m}} \right)^2 = n! \prod_{i=0}^{n-1} \left(\frac{m-i}{N-i} \right)^2 = n! \left(\frac{m}{N} \right)^{2n} e^{-o(n^{1/2})},$$

which clearly goes to infinity.

We will prove that

$$\frac{\mathbf{E}(X^2)}{\mathbf{E}(X)^2} \le f(\omega), \tag{1}$$

where $f(\omega) = e^{O(\omega^2)}$ for slowly growing ω.

The Paley-Zygmund inequality [12] implies that for $0 < \theta < 1$ we have

$$\mathbf{Pr}(X \ge \theta \mathbf{E}(X)) \ge (1-\theta)^2 \frac{\mathbf{E}(X)^2}{\mathbf{E}(X^2)}.$$

So, with $\theta = 1/2$ we can deduce that

$$\mathbf{Pr}(X > 0) \ge \frac{1}{4f(\omega)}.$$

Now let $p = \frac{2m}{N}$. Then we have, where $E_{n,p}$ is the set of edges of $D_{n,p}$,

$$\begin{aligned}
\mathbf{Pr}(X > 0 \text{ in } D_{n,p}) &\ge \mathbf{Pr}(X > 0 \text{ in } D_{n,p} \mid |E_{n,p}| \ge m)\mathbf{Pr}(|E_{n,p}| \ge m) \\
&\ge \sum_{m' \ge m} \mathbf{Pr}(X > 0 \text{ in } D_{n,m'}) - \mathbf{Pr}(|E_{n,p}| < m) \\
&\ge \frac{1}{4f(\omega)} - \mathbf{Pr}(|E_{n,p}| < m) \\
&\ge \frac{1}{4f(\omega)} - e^{-m/10} \\
&\ge \frac{1}{5f(\omega)}.
\end{aligned}$$

Now let $\omega_1 = \omega f(\omega)$ and let $m_1 = \omega_1 n^{3/2}$. Define p_1 by $p_1 = 1-(1-p)^{\omega_1}$ so that D_{n,p_1} can be expressed as the union of ω_1 independent copies of $D_{n,p}$. Note that $p_1 \le \omega_1 p$. Then we have

$$\mathbf{Pr}(X > 0 \text{ in } D_{n,p_1}) \ge \left(1 - \frac{1}{5f(\omega)}\right)^{\omega_1} = 1 - o(1).$$

The monotonicity of containing the square of a Hamilton cycle implies that D_{n,m_1} contains the square of a Hamilton cycle w.h.p. Here $m_1 = Np_1 \le \omega f(\omega) n^{3/2}$ and this clearly implies Theorem 1, after replacing ω by $\omega f(\omega)$.

To prove the result in $G_{n,p}$ we let $p_2 = 1-(1-p_1)^2$ and then we see that G_{n,p_1} is the graph underlying D_{n,p_2} and the square of a Hamilton cycle in D_{n,p_2} induces one in G_{n,p_1}. Finally, we obtain the result for $G_{n,m}$ by monotonicity.

3 Proof of (1)

Fix a square inducing permutation π, and let $H(\pi)$ be the square hamilton cycle induced by π. $H(\pi)$ consists of the red "distance 1 edges" $(\pi(i), \pi(i+1))$ and blue "distance 2 edges" $(\pi(i), \pi(i+2))$. We consider some other square inducing permutation $\widehat{\pi}$, and say that π shares a distance $k \in \{1,2\}$ edge with $\widehat{\pi}$ if $(\pi(i), \pi(i+k)) = (\widehat{\pi}(j), \widehat{\pi}(j+k))$ for some i, j. For F a set of distance 2 edges in $H(\pi) \cap H(\widehat{\pi})$ and F' in $H(\widehat{\pi})$ a set of paths ("paths" are all directed paths in this paper) of distance 1 edges in $\widehat{\pi}$, we say that F' *lies underneath* F if F is precisely the set of edges $(\widehat{\pi}(i), \widehat{\pi}(i+2))$ for indices i such that $(\widehat{\pi}(i), \widehat{\pi}(i+1))$ and $(\widehat{\pi}(i+1), \widehat{\pi}(i+2)$ are consecutive edges of a path in F'. A path in F' is a maximal sequence of the form $(\widehat{\pi}(l), \widehat{\pi}(l+1), \widehat{\pi}(l+2), \widehat{\pi}(l+3), \ldots) = (\pi(i), \pi(j), \pi(i+2), \pi(j+2), \ldots)$ that ends in $\pi(i+2s)$ or $\pi(j+2s)$ of vertices that are consecutive in $H(\widehat{\pi})$. These paths in F' may share endpoints. For example, a sequence $\pi(i), \pi(j), \pi(i+2), \pi(k), \pi(i+4)$ is considered to be two paths of length two if $k \neq j+2$. Note that every set F has a unique corresponding set of paths F' lying underneath F. Let $N^*(a,b,c,d)$ be the number of square inducing permutations $\widehat{\pi}$ such that:

(a) $H(\pi), H(\widehat{\pi})$ share b distance one edges (i.e. red edges), arranged in a common paths. These paths are obviously red.

(b) Let F be the set of shared (blue) distance 2 edges in $H(\pi) \cap H(\widehat{\pi})$, not including the shared (blue) distance 2 edges that arise from 2-paths of shared (red) distance 1 edges. That means not including edges of the form $(\pi(i), \pi(i+2))$ such that both $(\pi(i), \pi(i+1))$ and $(\pi(i+1), \pi(i+2))$ are shared edges. Then $|F| = d$ and the underlying set of paths F' has c paths and consequently $c+d$ edges.

We will first show that $N^*(a,b,c,d) \le N(a,b,c,d)$ for some explicit function $N(a,b,c,d)$. Note that we always have $0 \le a \le b \le n$ and $0 \le c \le d \le n$, which we assume below:

$$\frac{\mathbf{E}(X^2)}{\mathbf{E}(X)^2} \le \sum_b \sum_d \sum_a \sum_c \frac{n! N(a,b,c,d) \binom{N-2n+b}{m-2n+b}\binom{N-2n+d}{m-2n+d}/\binom{N}{m}^2}{\mathbf{E}(X)^2}$$
$$= \sum_b \sum_d \sum_a \sum_c \frac{N(a,b,c,d)\binom{N-2n+b}{m-n+b}\binom{N-2n+d}{m-2n+d}\binom{N}{m}^2}{n!\binom{N-n}{m-n}^4}. \tag{2}$$

Now observe that

$$\frac{\binom{N}{m}}{\binom{N-n}{m-n}} = \prod_{i=0}^{n-1} \frac{N-i}{m-i} = \left(\frac{N}{m}\right)^n \prod_{i=0}^{n-1} \left(1-\frac{i}{N}\right)\left(1-\frac{i}{m}\right)^{-1} \le \left(\frac{N}{m}\right)^n \exp\left\{\frac{n^2}{2m}\right\}$$

and

$$\frac{\binom{N-2n+b}{m-2n+b}}{\binom{N-n}{m-n}} = \prod_{i=0}^{n-b-1} \frac{m-(n+i)}{N-(n+i)} = \left(\frac{m}{N}\right)^{n-b} \prod_{i=0}^{n-b-1} \left(1-\frac{n+i}{m}\right)\left(1-\frac{n+i}{N}\right)^{-1}$$
$$\le_{\mathcal{O}} \left(\frac{m}{N}\right)^{n-b} \exp\left\{-\frac{3n^2-4nb}{2m}\right\},$$

where $f \le_{\mathcal{O}} g$ means $f = O(g)$. Consequently,

$$\frac{\binom{N-2n+b}{m-n+b}\binom{N-2n+d}{m-2n+d}\binom{N}{m}^2}{\binom{N-n}{m-n}^4} \le_{\mathcal{O}} \exp\left\{-\frac{2N}{m}\right\}\left(\frac{Ne^{2n/m}}{m}\right)^{b+d}.$$

Now we define an upper bound $N(a,b,c,d)$ for any $0 \le a \le b \le n$ and $0 \le c \le d \le n$. If $a = c = 0$ (implying $b = d = 0$), then we take $N(0,0,0,0) = n!$. Otherwise,

$$N(a,b,c,d) =$$
$$\frac{n}{a+c}\binom{b-1}{a-1}\binom{d-1}{c-1}\binom{a+c}{a}\times$$
$$\sum_{g_0=0}^{a+c}\binom{a+c}{g_0}\binom{n-b-c-d-1}{a+c-g_0-1} n^{a+2c-g_0}(n-a-b-2c-d+g_0)!,$$

where we use the convention $\binom{x}{-1} = 1$ for any $x > 0$. Here g_0 is the number of zero length gaps, see **S3** below.

We will now explain why $N^*(a,b,c,d) \le N(a,b,c,d)$. First recall that the number of integer solutions of

$$x_1 + \cdots + x_\ell = s \tag{3}$$

with $x_i \ge k$ (for each $1 \le i \le \ell$) is exactly $\binom{s-\ell(k-1)-1}{\ell-1}$. Obviously $N^*(0,0,0,0) \le n! = N(0,0,0,0)$. The remaining case we prove through the following steps:

S1 We choose a many positive lengths summing to b obtaining the lengths for paths in (a). This can be done in $\binom{b-1}{a-1}$ ways. (Apply (3) with $\ell = a$, $s = b$ and $k = 1$.)

S2 Similarly we choose c many lengths that are at least 2, summing to $c+d$ obtaining the lengths for underlying set of paths in (b). We have exactly $\binom{d-1}{c-1}$ choices. (Apply (3) with $\ell = c$, $s = c+d$ and $k = 2$.)

S3 These paths will have gaps (possibly zero length gaps) between. Let g_0 be the number of gaps of length zero. First we choose which gaps have zero length: $\binom{a+c}{g_0}$ ways. Note that it is actually not possible to have a zero gap between two paths (a), so we overcount here. It is possible to have a zero gap between two blue paths, or between a red and a blue path.

S4 Now we choose $a+c-g_0$ many positive lengths for gaps, summing to $n-b-c-d$. This gives $\binom{n-b-c-d-1}{a+c-g_0-1}$ many ways. (Apply (3) with $\ell = a+c-g_0$, $s = n-b-c-d$ and $k = 1$.)

S5 Now we specify the order of our red and blue paths in $\widehat{\pi}$. We already have an ordered set of a many lengths of red paths, and another ordered set of c lengths of blue paths. Thus the number of ways to specify an ordering of these $a+c$ many paths together is $\binom{a+c}{a}$.

S6 We choose a "starting index" (n choices) for the first of our $a+c$ many paths. Once this choice is made we have specified the positions of all the paths and gaps in $\widehat{\pi}$. But each possible configuration is counted $a+c$ many times. So for this part we multiply by $\frac{n}{a+c}$.

Let us describe what we know about $\widehat{\pi}$ at this point. By "starting index" we mean we choose an index i such that the first of our $a+c$ paths, if it has say length L, will be the path $\widehat{\pi}(i), \widehat{\pi}(i+1), \ldots, \widehat{\pi}(i+L)$. Then the first gap, say if it has length L' will be $\widehat{\pi}(i+L), \ldots, \widehat{\pi}(i+L+L')$. Note that we have not yet specified any of the actual values of $\widehat{\pi}(j)$ for any j. We will start doing that next.

S7 We choose the value that $\widehat{\pi}$ takes at the first index of each path in (a) and each underlying path in (b), except for those paths which come immediately after a gap of length zero. Since there are g_0 zero length gaps we have n^{a+c-g_0} choices. Now we choose the second vertex for each underlying paths having n^c choices.

For example if we already know a certain path (which does not come immediately after a zero length gap) in (a) is $\widehat{\pi}(j), \ldots, \widehat{\pi}(j+L)$ then we choose the value of $\widehat{\pi}(j)$. Note that there is some index j' such that $\widehat{\pi}(j) = \pi(j')$, and since this is a path of shared edges we must have $\widehat{\pi}(j+t) = \pi(j'+t)$ for $0 \le t \le L$.

Similarly if we already know a certain path (which does not come after a zero length gap) in (b) is $\widehat{\pi}(j), \ldots, \widehat{\pi}(j+L)$ then we choose the value of $\widehat{\pi}(j), \widehat{\pi}(j+1)$. Note that there are some indices j', j'' such that $\widehat{\pi}(j) = \pi(j'), \widehat{\pi}(j+1) = \pi(j'')$, and so $\widehat{\pi}(j), \widehat{\pi}(j+1), \widehat{\pi}(j+2), \widehat{\pi}(j+3) \ldots = \pi(j'), \pi(j''), \pi(j'+2), \pi(j''+2) \ldots$.

For paths of type (a) or (b) that do come immediately after a zero gap, the value of $\widehat{\pi}$ at the first index is already determined by the path that came before. However we always choose the value of $\widehat{\pi}$ at the second index of each path in (b).

S8 Finally we choose the values of $\widehat{\pi}$ at all of the indices that are not in any of the paths from (a) and (b). There are $a+b+2c+d-g_0$ many vertices on these paths.

Observe that in (2) the first term ($b = d = a = c = 0$) is negligible. Thus, from now on we assume that $a + c \geq 1$ and get

$$\begin{aligned}\frac{\mathbf{E}(X^2)}{\mathbf{E}(X)^2} &\leq \sum_b \sum_d \sum_a \sum_c \frac{N(a,b,c,d)\binom{N-2n+b}{m-n+b}\binom{N-2n+d}{m-2n+d}\binom{N}{m}^2}{n!\binom{N-n}{m-n}^4} \\ &\leq_{\mathcal{O}} \frac{1}{n!}\exp\left\{-\frac{2N}{m}\right\}\sum_b\sum_d\sum_a\sum_c\sum_{g_0}\frac{n}{a+c}\binom{b-1}{a-1}\binom{d-1}{c-1}\binom{a+c}{a}\binom{a+c}{g_0} \\ &\quad\times\binom{n-b-c-d-1}{a+c-g_0-1}n^{a+2c-g_0}(n-a-b-2c-d+g_0)!\left(\frac{Ne^{2n/m}}{m}\right)^{b+d}. \quad (4)\end{aligned}$$

Now let $\delta = \omega/n^{1/2}$ and consider two cases.

Case 1: $b \leq \delta n$ **and** $d \leq \delta n$

Since $a \leq b$ and $c \leq d$, we have $a, c \leq \delta n$ and $g_0 \leq a + c \leq 2\delta n$. Therefore,

$$\begin{aligned}\frac{n^{a+2c-g_0}(n-a-b-2c-d+g_0)!}{(n-b-d)!} &= \frac{n^{a+2c-g_0}}{(n-b-d)_{a+2c-g_0}} \leq \frac{n^{a+2c-g_0}}{(n-5\delta n)^{a+2c-g_0}} \\ &= \left(\frac{1}{1-5\delta}\right)^{a+2c-g_0} \leq \left(\frac{1}{1-5\delta}\right)^{3\delta n} \leq e^{20\delta^2 n} = e^{20\omega^2}\end{aligned}$$

and so (4) is at most

$$\begin{aligned}&e^{20\omega^2}\frac{1}{n!}\exp\left\{-\frac{2N}{m}\right\}\sum_b\sum_d\sum_a\sum_c\sum_{g_0}\frac{n}{a+c}\binom{b-1}{a-1}\binom{d-1}{c-1}\binom{a+c}{a}\binom{a+c}{g_0} \\ &\quad\times\binom{n-b-c-d-1}{a+c-g_0-1}(n-b-d)!\left(\frac{Ne^{2n/m}}{m}\right)^{b+d}.\end{aligned}$$

Now we use the Vandermonde identity to obtain

$$\sum_{g_0=0}^{a+c}\binom{a+c}{g_0}\binom{n-b-c-d-1}{a+c-g_0-1} = \binom{n+a-b-d-1}{a+c-1}$$

to get the bound

$$\begin{aligned}&e^{20\omega^2}\frac{1}{n!}\exp\left\{-\frac{2N}{m}\right\}\sum_b\sum_d\sum_a\sum_c\frac{n}{a+c}\binom{b-1}{a-1}\binom{d-1}{c-1}\binom{a+c}{a} \\ &\quad\times\binom{n+a-b-d-1}{a+c-1}(n-b-d)!\left(\frac{Ne^{2n/m}}{m}\right)^{b+d} \\ &\qquad\leq_{\mathcal{O}} e^{20\omega^2}\frac{1}{n!}\exp\left\{-\frac{2N}{m}\right\}\sum_b\sum_d\sum_a\sum_c\binom{b-1}{a-1}\binom{d-1}{c-1}\binom{a+c}{a} \\ &\qquad\quad\times\binom{n+a-b-d-1}{a+c}(n-b-d)!\left(\frac{Ne^{2n/m}}{m}\right)^{b+d}. \quad (5)\end{aligned}$$

Now note that

$$\sum_{c=0}^{d}\binom{d-1}{c-1}\binom{a+c}{a}\binom{n+a-b-d-1}{a+c}=\binom{n+a-b-d-1}{a}\sum_{c}\binom{d-1}{d-c}\binom{n-b-d-1}{c}$$
$$=\binom{n+a-b-d-1}{a}\binom{n-b-2}{d}.$$

Hence, (5) can be bounded by

$$e^{20\omega^2}\frac{1}{n!}\exp\left\{-\frac{2N}{m}\right\}\sum_{b}\sum_{d}\sum_{a}\binom{b-1}{a-1}\binom{n+a-b-d-1}{a}\binom{n-b}{d}(n-b-d)!\left(\frac{Ne^{2n/m}}{m}\right)^{b+d}$$
$$\le e^{20\omega^2}\frac{1}{n!}\exp\left\{-\frac{2N}{m}\right\}\sum_{b}\sum_{d}\sum_{a}\binom{b-1}{a-1}\binom{n-d-1}{a}\binom{n-b}{d}(n-b-d)!\left(\frac{Ne^{2n/m}}{m}\right)^{b+d}$$
$$=e^{20\omega^2}\frac{1}{n!}\exp\left\{-\frac{2N}{m}\right\}\sum_{b}\sum_{d}\binom{n+b-d-2}{b}\binom{n-b}{d}(n-b-d)!\left(\frac{Ne^{2n/m}}{m}\right)^{b+d}\qquad(6)$$

and note that since

$$\frac{(n+b-d-2)_b(n-b)_d(n-b-d)!}{n!}=\frac{(n+b-d-2)_b}{(n)_b}\le\left(\frac{n+b}{n-b}\right)^b\le\left(\frac{n+\delta n}{n-\delta n}\right)^{\delta n}$$
$$=\left(\frac{1+\delta}{1-\delta}\right)^{\delta n}\le(1+3\delta)^{\delta n}\le e^{3\omega^2}$$

we can say that (6) is at most

$$e^{23\omega^2}\exp\left\{-\frac{2N}{m}\right\}\sum_{b}\sum_{d}\frac{\left(\frac{Ne^{2n/m}}{m}\right)^{b+d}}{b!d!}\le e^{23\omega^2}\exp\left\{-\frac{2N}{m}\right\}\sum_{b=0}^{\infty}\frac{\left(\frac{Ne^{2n/m}}{m}\right)^{b}}{b!}\sum_{d=0}^{\infty}\frac{\left(\frac{Ne^{2n/m}}{m}\right)^{d}}{d!}$$
$$=e^{23\omega^2}\exp\left\{-\frac{2N}{m}\right\}\exp\left\{\frac{Ne^{2n/m}}{m}\right\}\exp\left\{\frac{Ne^{2n/m}}{m}\right\}$$
$$=e^{23\omega^2}\exp\left\{\frac{2N}{m}\left(e^{2n/m}-1\right)\right\}=e^{23\omega^2}\exp\left\{O\left(\frac{Nn}{m^2}\right)\right\}\le e^{25\omega^2}.$$

Case 2: $\max\{b,d\}\ge\delta n$

We will show that the term of (4)

$$\frac{1}{n!}\exp\left\{-\frac{2N}{m}\right\}\sum_{a}\sum_{c}\sum_{g_0}\frac{n}{a+c}\binom{b-1}{a-1}\binom{d-1}{c-1}\binom{a+c}{a}\binom{a+c}{g_0}$$
$$\times\binom{n-b-c-d-1}{a+c-g_0-1}n^{a+2c-g_0}(n-a-b-2c-d+g_0)!\left(\frac{Ne^{2n/m}}{m}\right)^{b+d}\qquad(7)$$

is very small. First note that

$$\frac{n^{a+2c-g_0}(n-a-b-2c-d+g_0)!}{n!}$$
$$\leq_{\mathcal{O}} \frac{n^{a+2c-g_0}\sqrt{n-a-b-2c-d+g_0}\left(\frac{n-a-b-2c-d+g_0}{e}\right)^{n-a-b-2c-d+g_0}}{\sqrt{n}\left(\frac{n}{e}\right)^n} \leq e^{O(b+d)}n^{-b-d}.$$

Thus, we have that (7) is at most

$$\begin{aligned}
&e^{O(b+d)}\left(\frac{N}{mn}\right)^{b+d}\sum_a\sum_c\sum_{g_0}\frac{n}{a+c}\binom{b-1}{a-1}\binom{d-1}{c-1}\binom{a+c}{a}\binom{a+c}{g_0}\binom{n-b-c-d-1}{a+c-g_0-1}\\
&=e^{O(b+d)}\left(\frac{N}{mn}\right)^{b+d}\sum_a\sum_c\frac{n}{a+c}\binom{b-1}{a-1}\binom{d-1}{c-1}\binom{a+c}{a}\binom{n+a-b-d-1}{a+c-1}\\
&=e^{O(b+d)}\left(\frac{N}{mn}\right)^{b+d}\sum_a\sum_c\binom{b-1}{a-1}\binom{d-1}{c-1}\binom{a+c}{a}\binom{n+a-b-d}{a+c}\\
&\leq e^{O(b+d)}\left(\frac{N}{mn}\right)^{b+d}\sum_a\sum_c\binom{b-1}{a-1}\binom{d-1}{c-1}\binom{a+c}{a}\binom{n-d}{a+c}\\
&=e^{O(b+d)}\left(\frac{N}{mn}\right)^{b+d}\sum_a\sum_c\binom{b-1}{a-1}\binom{d-1}{c-1}\binom{n-d-a}{c}\binom{n-d}{a}\\
&=e^{O(b+d)}\left(\frac{N}{mn}\right)^{b+d}\sum_a\binom{b-1}{a-1}\binom{n-a-1}{d}\binom{n-d}{a}\\
&\leq e^{O(b+d)}\left(\frac{N}{mn}\right)^{b+d}\sum_a\binom{b-1}{a-1}\binom{n}{d}\binom{n-d}{a}\\
&=e^{O(b+d)}\left(\frac{N}{mn}\right)^{b+d}\binom{n}{d}\binom{n+b-d-1}{b}\\
&\leq e^{O(b+d)}\left(\frac{N}{m}\right)^{b+d}\frac{1}{b!d!}\\
&\leq \left(\frac{NO(1)}{mb}\right)^b\left(\frac{NO(1)}{md}\right)^d
\end{aligned}$$

which is exponentially small. To see this, first note that the final expression is symmetric. Hence, without loss of generality we can assume that $b \geq \delta n$. Then

$$\left(\frac{NO(1)}{mb}\right)^b = \left(\frac{O(1)}{\omega^2}\right)^b = e^{-\Omega(b\log\omega)}$$

and since the function $f(x) = (C/x)^x$ is maximized at $x = C/e$ where we have $f(x) = e^{C/e}$, we may bound

$$\left(\frac{NO(1)}{md}\right)^d = e^{O(N/m)} = e^{O(b)}.$$

This completes the proof of (1) and hence Theorem 1.

4 Proof of Theorem 2

4.1 Preliminaries

It will be convenient for the computations to assume that the edges X will be given as $X = X_1 \cup X_2$ where each of the sets in this partition are independent random subsets of $E(K_n)$ where each edge is independently included with probability $p = \frac{K \log^{1/3} n}{n^{2/3}}$.

Assume first that $n = 2m$ is even. It follows from Erdős and Rényi [5] that w.h.p. the edges X_1 contain a perfect matching M. By symmetry, it will be a random matching of K_n that is independent of G. It can therefore be derived from a random permutation $\pi = (z_1, z_2, \ldots, z_n)$ via $M = \{e_1, e_2, \ldots, e_m\}$ where $e_i = \{z_{2i-1}, z_{2i}\}, i = 1, 2, \ldots, m$.

Now define a graph Γ with vertex set M and an edge $\{e, f\}, e, f \in M$ whenever the subgraph $H_{e,f}$ of $G + X_1$ induced by the four vertices in $e \cup f$ is K_4. We argue that

$$\text{W.h.p., } \Gamma \text{ has minimum degree at least } \beta_1 n \text{ where } \beta_1 = (2\alpha - 1)^2/3. \tag{8}$$

To see this consider a fixed edge $e = \{x, y\} \in M$. Let $N(a)$ denote the set of neighbors of vertex a in G and note that $|N(x) \cap N(y)| \geq (2\alpha - 1)n$. The probability that another edge $\{u, v\} \in M$ satisfies $u, v \in N(x) \cap N(y)$ is at least $(1 - o(1))(2\alpha - 1)^2/2$. Thus the degree d_e of edge e has expectation at least $(1-o(1))(2\alpha-1)n/2$ in Γ. Swapping a pair in permutation π can only change d_e by at most two. Applying a version of the Azuma-Hoeffding inequality (see for example McDiarmid [10] or Frieze and Pittel [6]) we see that $\mathbf{Pr}(d_e \leq (2\alpha - 1)^2 n/3) \leq e^{-\Omega((2\alpha-1)^4 n)}$. This verifies (8), after inflating the probability bound by m. Note that only the $G + X_1$-edges of Γ are used here.

We prove next the following lemma.

Lemma 3 *Γ is connected w.h.p.*

Proof It follows from (8) that we only need to show there are no components S of size $s \in [\beta_1 n/2, m/2]$. But then

$$\begin{aligned}
\mathbf{Pr}(\exists \text{ component of size } s \in [\beta_1 n/2, m/2]) &\leq \sum_{s=\beta_1 n/2}^{m/2} \binom{n}{s} (1-p^3)^{s(\alpha n - 2s)/2} \qquad (9)\\
&\leq \sum_{s=\beta_1 n/2}^{n/4} \left(\frac{ne}{s}\right)^s e^{-s(\alpha n - 2s)p^3/2}\\
&\leq \sum_{s=\beta_1 n/2}^{n/4} \left(\frac{ne}{s} \cdot e^{-\frac{\beta_1}{4} K^3 (\alpha - \frac{1}{2}) \log n}\right)^s\\
&= o(1).
\end{aligned}$$

Explanation of (9)**:** For each $e \in S$ there are at least $(\alpha n - 2s)$ vertices T outside $\bigcup_{e \in S} e$. For each vertex $w \in T$ we have a matching edge $f \in \bar{S}$ such that e and f are joined by an edge

from G. The term p^3 accounts for the probability that X_2 will provide another three edges to create a K_4. We divide by two in $s(\alpha n - 2s)/2$ to account for there being two choices for $w \in f$. □

4.2 2-paths

A *2-path* is a sequence of vertices $(x_1, x_2, \ldots, x_{2k})$ such that (i) $(x_1, x_2, \ldots, x_{2k})$ is a path in $G + X$, (ii) $\{x_{2i-1}, x_{2i}\} \in M$, $i = 1, 2, \ldots, k$, and (iii) $\{x_i, x_{i+2}\}$ are edges of $G + X$ for $i = 1, 2, \ldots, 2k - 2$ (see Figure 1).



Figure 1: A 2-path for $k = 7$.

In a 2-path we refer to the edges $\{x_{2i-1}, x_{2i}\}, i = 1, 2, \ldots, k$ as the *pillars*.

We now define a *rotation* with $\{x_1, x_2\}$ as the *fixed end* and $\{x_{2k-1}, x_{2k}\}$ as the *rotated end*. Suppose that for some $\ell \leq k-2$ we have that $\{x_{2\ell-1}, x_{2k}\}, \{x_{2\ell}, x_{2k-1}\}, \{x_{2\ell}, x_{2k}\}$ are all edges of $G + X$. Then we obtain a new 2-path $(x_1, x_2, \ldots, x_{2\ell}, x_{2k}, x_{2k-1}, \ldots, x_{2\ell+1})$ (see Figure 2).

4.3 Algorithm ERA

Extension-Rotation algorithm
The algorithm begins by choosing an arbirtrary edge $e \in M$ and letting path $P_1 = e$.

Basic Idea It proceeds in rounds. At the beginning of round k we will have a 2-path $P_k =$



Figure 2: A rotated path for $k = 7$ and $\ell = 3$.

$(x_1, x_2, \ldots, x_{2k})$. A round consists of the following: Let $Q_0 = P_k$ and then for $i = 1, 2, \ldots$, if necessary, grow a set of paths $Q_1, Q_2, \ldots$. Each Q_i is obtained from some $Q_j, j < i$ by a single rotation.

We continue until either we make a simple extension (defined below) or we make a cycle extension (defined below) or fail.

Simple Extensions The process is curtailed if at any point the procedure generates a path $P = (y_1, y_2, \ldots, y_{2k})$ and an edge $\{u, v\} \in M$ disjoint from $V(P)$ such that $(y_1, y_2, \ldots, y_{2k}, u, v)$ is a 2-path. In which case we can extend our current 2-path to one of length $2k + 2$ and end the round. We call this a *simple extension*.

Cycle extensions If we do not find a simple extension, then we see if there is a path $P = (y_1, y_2, \ldots, y_{2k}) \in \mathcal{P}_L$ such that $G + X$ contains the path $(y_{2k-1}, y_1, y_{2k}, y_2)$ (see Figure 3).



Figure 3: Closing a 2-path with $(y_{2k-1}, y_1, y_{2k}, y_2)$.

We say that we *close the path* to create a cycle $C = (y_1, y_2, \ldots, y_{2k}, y_1)$. If we find such an edge and $k = n/2$ then we have found the square of a Hamilton cycle. Otherwise, we seek a *cycle extension*. By this we mean that find an edge $\{u, v\} \in M$ disjoint from $V(P)$ such that and $1 \le \ell < k$ such that $G + X$ contains the path $(y_{2\ell-1}, u, y_{2\ell}, v)$. In which case we now have the 2-path $(y_{2\ell+1}, y_{2\ell+2}, \ldots, y_{2k}, y_{2k-1}, y_1, y_2, \ldots, y_{2\ell-1}, y_{2\ell}, u, v)$. We call this a *cycle extension* (see Figure 4). If no such pair $\ell, \{u, v\}$ exists then we fail.

We need to be a little more precise about the order of rotations.

Step 1 We start a round with $P_k = (x_1, x_2, \ldots, x_{2k})$. We then do a set of rotations with $e = \{x_1, x_2\}$ as the fixed end, one for each neighbor of $\{x_{2k-1}, x_{2k}\}$ in Γ. We will use the edges of $G + X_1$ when we check for simple extensions. Assuming there are no simple extensions we generate a set of 2-paths $Q_1, Q_2, \ldots, Q_L, L \ge \beta_1 n$. The end pillar of P_i, other than $\{x_1, x_2\}$, will be denoted by e_i for $i = 1, 2, \ldots, L$.

Step 2 After this, we take each Q_i in turn and do a set of rotations with e_i as the fixed end and e as the rotated end, using the edges of $G + X_1$ for this purpose.

Step 3 If we fail to obtain a simple extension, then we use the $G + X$ edges to look for a cycle extension, using all of the 2-paths generated for this task.



Figure 4: A cycle extension.

4.4 Analysis of ERA

Lemma 4 *W.h.p. algorithm* ERA *succeeds in finding the square of a Hamilton cycle.*

Proof We argue that w.h.p. we can always find an X_2 edge to close a path if there is no simple extension. For a fixed round we have from (8) that

$$\mathbf{Pr}(\text{No path of } \mathcal{P}_k \text{ can be closed}) \leq (1-p^3)^{\beta_1^2 n^2/2} \leq e^{-\frac{1}{2}\beta_1^2 K^3 \log n}. \tag{10}$$

Since there at most $n/2$ rounds we see that w.h.p. there is at least one path in a round that can be closed, if needed.

Having closed a 2-path, the existence of $\ell, \{u, v\}$ follows from the connectivity of Γ, see Lemma 3. □

When $n = 2m+1$ is odd, we use extra $Kn^{4/3}\log^{1/3} n$ edges X_3 (chosen independently from X_1 and X_2) to find a subgraph as in Figure 5.



Figure 5: A graph used for n odd.

We can then use a, d and b, e as pillars and basically proceed as in the even case, making sure to avoid breaking up this subgraph and follow its vertices as (d, a, c, b, e).

4.5 A lower bound

It is as well to consider lower bounds on the number of random edges needs to add to a graph G to obtain the kth power of Hamilton cycle. For this we consider the complete bipartite graph $G = K_{s,t}$ with bipartition A, B and where $s = |A| = \alpha n$ and $s + t = n$. We will be thinking here of the case where α is a small constant and so it does not fit exactly into the assumptions of Theorem 2. In $K_{s,t}$ the lower bound is much less than $n/2$ and have no lower bounds for the case where the minimum degree significantly exceeds $n/2$.

We can associate a sequence σ_H of length n over the alphabet $\{A, B\}$ with a Hamilton cycle H in $K_{s,t}$. The ith symbol will be an A if and only if the ith vertex of the cycle is in A. Only AB edges are in G and it is not difficult to show by examining σ_H that at most $2ks$ of the edges of H can be of this type. It follows that if we add edges to G with probability p then the expected number of kth powers will be at most $n!p^{k(n-2s)}$. Thus we require $p \geq n^{-1/k(1-2\alpha)}$ or at least $n^{2-1/k(1-2\alpha)}$ random edges. In particular, for $k = 2$ this implies that we need $n^{2-1/(2-4\alpha)}$ random edges, which for small α yields $n^{3/2-O(\alpha)}$. This is quite optimal, since due to Theorem 1 $\omega n^{3/2}$ is the trivial upper bound.

5 Final Remarks

It follows from Theorem 1, a result of Riordan [14] and the embedding theorem in Dudek, Frieze, Ruciński and Šileikis [4] that w.h.p. the random r-regular $G_{n,r}$ contains the kth power of a Hamilton cycle as long as $n^{1-1/k} \ll r \ll n$, $k \geq 2$.

It might be also of some interest to extend Theorem 2 for any $k \geq 3$ and determine how many random edges need to be added to a graph of order n with minimum degree αn in order that it contains the kth power of a Hamilton cycle w.h.p.

6 Acknowledgments

It is a pleasure to thank Asaf Ferber and Matthew Kwan for a close reading of an earlier version of the paper. They pointed to an error in the proof of Theorem 1 that led us to switch from $G_{n,p}$ to $D_{n,m}$.

References

[1] M. Ajtai, J. Komlós and E. Szemerédi, *The first occurrence of Hamilton cycles in random graphs*, Annals of Discrete Mathematics **27** (1985), 173–178.

[2] T. Bohman, A.M. Frieze and R. Martin, *How many random edges make a dense graph Hamiltonian?*, Random Structures and Algorithms 22 (2003) 33-42.

[3] B. Bollobás, *The evolution of sparse graphs*, in Graph Theory and Combinatorics, Academic Press, Proceedings of Cambridge Combinatorics, Conference in Honour of Paul Erdős (B. Bollobás; Ed) (1984), 35–57.

[4] A. Dudek, A.M. Frieze, A. Ruciński and M. Šileikis, *Embedding the Erdős-Rényi hypergraph into the random regular hypergraph and Hamiltonicity*, Journal of Combinatorial Theory, Ser. B **122** (2017), 719–740.

[5] P. Erdős and A. Rényi, *On the existence of a factor of degree one of a connected random graph*, Acta. Math. Acad. Sci. Hungar. 17 (1966) 359-368.

[6] A.M. Frieze and B. Pittel, Perfect matchings in random graphs with prescribed minimal degree, *Trends in Mathematics, Birkhauser Verlag, Basel* (2004) 95-132.

[7] J. Komlós and E. Szemerédi, *Limit distributions for the existence of Hamilton circuits in a random graph*, Discrete Mathematics **43** (1983), 55–63.

[8] J. Komlós, N. Sarközy and E. Szemerédi, *Proof of the Seymour conjecture for large graphs*, Annals of

[9] D. Kühn and D. Osthus, *On Pósa's conjecture for random graphs*, SIAM Journal on Discrete Mathematics **26** (2012), 1440–1457.

[10] C. McDiarmid, On the method of bounded differences, *in Surveys in Combinatorics, ed. J. Siemons, London Mathematical Society Lecture Notes Series* 141, Cambridge University Press, 1989.

[11] R. Nenadov and N. Škorić, *Powers of cycles in random graphs and hypergraphs*, arXiv:1601.04034v1.

[12] R. Paley and A. Zygmund, *A note on analytic functions in the unit circle*, Proceedings of the Cambridge Philosophical Society **28** (1932), 266-272

[13] L. Pósa, *Hamiltonian circuits in random graphs*, Discrete Mathematics **14** (1976), 359–364.

[14] O. Riordan, *Spanning subgraphs of random graphs*, Combinatorics, Probability and Computing **9** (2000), 125–148.